

# Normed convergence property for hypergroups admitting an invariant measure

C. R. E. Raja

## Abstract

We prove that a hypergroup admitting a countable basis and an invariant Haar measure has normed convergence property if and only if it is compact.

*AMS Mathematics Subject Classification 2000:* 43A62, 60B15.

*Keywords:* Hypergroups, probability measures, weak topology and invariant measure.

Let  $K$  be a (locally compact) hypergroup and  $\mathcal{P}(K)$  be the space of all regular Borel probability measures on  $K$  equipped with the weak topology. For  $x \in K$ ,  $\delta_x$  denotes the probability measure concentrated at  $x$  and for  $\mu \in \mathcal{M}^1(K)$ ,  $x\mu$  and  $\mu x$  denote  $\delta_x * \mu$  and  $\mu * \delta_x$  respectively. Also, for  $\mu_1, \mu_2 \in \mathcal{P}(K)$ ,  $\mu_1\mu_2$  denotes  $\mu_1 * \mu_2$ .

**Definition** *A hypergroup  $K$  is said to have normed convergence property if for any sequence  $(\mu_n)$  in  $\mathcal{P}(K)$ , there exists a sequence  $(x_n)$  in  $K$  such that  $(x_n\nu_n)$  is relatively compact where  $\nu_n = \mu_n \cdots \mu_2\mu_1$  for all  $n \geq 1$ .*

It is well-known that a locally compact group has normed convergence property if and only if it is compact (see Theorem 2.3.2 of [5]) and a Moore hypergroup has normed convergence property if and only if it is compact (see [3]). In this short note we extend the afore-mentioned result to hypergroups admitting an invariant Haar measure.

Let  $K$  be a hypergroup admitting a right-invariant Haar measure  $m$ . Let  $L^p(K)$  be the space of  $p$ -integrable functions on  $K$  for  $1 \leq p < \infty$ . Let  $\mathcal{M}^1(K)$  be the space of all non-negative measures on  $K$  with total measure

less than are equal to 1. Now, for any  $\mu \in \mathcal{M}^1(K)$  and for  $1 \leq p \leq \infty$ , define  $P_\mu: L^p(K) \rightarrow L^p(K)$  by

$$P_\mu(f)(x) = \int f(xy) d\mu(y) = \int f(y) d(x\mu)(y)$$

where  $f(xy) = \int f(z) d(\delta_x * \delta_y)(z)$ , for all  $x \in K$ . It is easy to see that  $P_\mu$  is a contraction on  $L^p(K)$  for any  $\mu \in \mathcal{M}^1(K)$  and any  $1 \leq p \leq \infty$ . The map  $\mu \mapsto P_\mu$  is an injection, so we may view  $\mathcal{M}^1(K)$  as a subspace of bounded linear operators on  $L^2(K)$ .

We now recall the following topologies on  $\mathcal{M}^1(K)$ :

1. weak topology on  $\mathcal{M}^1(K)$  is the weakest topology for which the function  $\mu \mapsto \mu(f)$  is continuous for all bounded continuous functions  $f$  on  $K$ ;
2. vague topology on  $\mathcal{M}^1(K)$  is the weakest topology for which the function  $\mu \mapsto \mu(f)$  is continuous for all continuous functions  $f$  with compact support;
3. weak operator topology (respectively, strong operator topology) on  $\mathcal{M}^1(K)$  is the subspace topology inherited from the space of bounded linear operators on  $L^2(K)$  with weak operator topology (respectively, strong operator topology).

For a locally compact group  $G$ , it is well-known that on  $\mathcal{M}^1(G)$  the vague and weak operator topologies coincide and on  $\mathcal{P}(G)$ , the weak, vague and weak operator and strong operator topologies coincide (see Lemma 6.1.23 of [5]). The following extension of this result may be proved arguing as in Lemma 6.1.23 of [5] and we omit the details.

**Proposition 1** *Let  $K$  be a hypergroup admitting a countable basis and a right-invariant measure. Then we have the following:*

1. *on  $\mathcal{M}^1(K)$ , the vague and weak operator topologies coincide;*
2. *on  $\mathcal{P}(K)$ , the weak, vague and weak operator and strong operator topologies coincide.*

*In particular,  $\mathcal{M}^1(K)$  is compact in the weak operator topology.*

We next prove the following:

**Lemma 1** *Let  $K$  be as in Proposition 1. Suppose  $(\mu_n)$  is a sequence in  $\mathcal{P}(K)$  such that  $(x_n\mu_n)$  is relatively compact. Then there exists a  $f \in L^2(K)$  such that  $\|P_{\mu_n}(f)\|_2 \not\rightarrow 0$ .*

**Proof** Since  $(x_n\mu_n)$  is relatively compact, there exists a  $\delta > 0$  and a compact set  $C$  such that  $x_n\mu_n(C) > \delta$  for all  $n \geq 1$ .

Let  $f$  be a continuous function on  $K$  with compact support such that  $0 \leq f \leq 1$  and  $f(x) = 1$  for all  $x \in C$ . Then

$$P_{\mu_n}(f)(x_n) = \int f(y)d(x_n\mu_n)(y) > \delta$$

for  $n \geq 1$ . Let  $(h_j)$  be a compactly supported left approximate identity in the space of continuous functions vanishing at infinity. Then as in Proposition 3.1 of [4], we have

$$\|P_{\mu_n}(f)\|_\infty \leq \|h_j * f - f\|_\infty + \|P_{\mu_n}(f)\|_2 \|h_j\|_2$$

where  $h_j * f(y) = \int h_j(x)f(xy)dm(x)$ . Since  $\|P_{\mu_n}(f)\|_\infty > \delta > 0$  for all  $n \geq 1$ , we get that  $\|P_{\mu_n}(f)\|_2 \not\rightarrow 0$ .

**Theorem 1** *Let  $K$  be a hypergroup admitting a right-invariant Haar measure and a countable basis. Then  $K$  has normed convergence property if and only if  $K$  is compact.*

**Proof** It is well-known that compact hypergroups have normed convergence property. We now prove the 'only if' part. Suppose  $K$  has normed convergence property. Let  $\mu \in \mathcal{P}(K)$  be a symmetric measure such that the support of  $\mu$  is  $K$ . Then by assumption, there exists a sequence  $(x_n)$  such that  $(x_n\mu^n)$  is relatively compact. By Lemma 1, we get that

$$\|P_\mu^n(f)\|_2 \not\rightarrow 0$$

as  $n \rightarrow \infty$  for some continuous function  $f$  with compact support. By convergence of alternating sequences (see [1]), in the strong operator topology

$$P_\mu^{2n} \rightarrow P$$

where  $P$  is a bounded linear operator on  $L^2(K)$ . This implies that  $P$  is a self-adjoint projection. By Proposition 1, there exists a  $\rho \in \mathcal{M}^1(K)$  such that  $P = P_\rho$ . Since  $P_\mu^n(f) \not\rightarrow 0$ ,  $\rho \neq 0$  and  $\rho^2 = \rho$ . This means that  $\rho$  is the normalized Haar measure on a compact subhypergroup  $H$  of  $K$  (see [2]). Since  $P_\mu^2 P = P$ , we get that the support of  $\mu^2$  is contained in  $H$  (see [2]). Since the support of  $\mu$  is  $K$ , we get that  $K = H$  and hence  $K$  is compact.

## References

- [1] Bela, Sz.-Nagy and Ciprian Foias, *Harmonic analysis of operators on Hilbert space*, North-Holland, 1970.
- [2] W. R. Bloom and H. Heyer, The Fourier transform for probability measures on hypergroups, *Rendiconti di Matematica* (7) 2, 315-334 (1982).
- [3] W. R. Bloom and H. Heyer, Convergence of convolution products of probability measures on hypergroups, *Rendiconti di Matematica* (7) 2, 547-563 (1982).
- [4] Y. Derriennic and M. Lin, Convergence of iterates of averages of certain operator representations and of convolution powers, *Journal of Functional Analysis* 85, 86-102 (1989).
- [5] H. Heyer, *Probability measures on locally compact groups*, Springer-Verlag, 1977.

C. Robinson Edward Raja,  
Stat-Math Unit,  
Indian Statistical Institute,  
8th Mile Mysore Road,  
R. V. College Post,  
Bangalore - 560 059.  
India. e-mail: creraja@isibang.ac.in  
or  
e-mail: raja\_robinson@hotmail.com